

The AP Physics C: Mechanics equation sheet, annotated

Given in both sections (College Board, 2025): $G = 6.67 \times 10^{-11} \text{ N}\cdot\text{m}^2/\text{kg}^2$; $g = 9.8 \text{ m/s}^2 = 9.8 \text{ N/kg}$; prefixes tera to pico; units Hz, J, kg, m, N, s, W; sin, cos, tan for 0° , 30° , 37° , 45° , 53° , 60° , 90° (37° and 53° are the 3-4-5 triangle). Conventions: inertial frame, no air resistance, ideal springs and strings unless stated.

Translational	Use when
$v_x = v_{x0} + a_x t$; $x = x_0 + v_{x0} t + \frac{1}{2} a_x t^2$; $v_x^2 = v_{x0}^2 + 2a_x(x - x_0)$	constant acceleration only
$\Delta x = \int v_x dt$; $\Delta v_x = \int a_x dt$	$v(t)$ or $a(t)$ varies: integrate (area under the graph)
$x_{\text{cm}} = \sum m_i x_i / \sum m_i$; $r_{\text{cm}} = \int r dm / \int dm$; $\lambda = dm/d\ell$	centre of mass of points; of a continuous body; non-uniform rods ($dm = \lambda d\ell$)
$a_{\text{sys}} = \Sigma F / m_{\text{sys}} = F_{\text{net}} / m_{\text{sys}}$	Newton's second law for the system
$ F_g = Gm_1 m_2 / r^2$; $ F_f \leq \mu F_N $; $F_s = -k\Delta x$	gravitation (centre to centre); friction ($\leq$ static, = kinetic); springs (restoring)
$a_c = v^2/r = r\omega^2$; $T = 1/f$	circular motion
$K = \frac{1}{2}mv^2$; $W = \int F \cdot dr$; $\Delta K = \Sigma W_i = \Sigma F_{i,j} \cdot d_i$	energy; work by a variable force; the work-energy theorem
$\Delta U = -\int F_{cf} \cdot dr$; $F_x = -dU/dx$	U from a conservative force; force as minus the slope of $U(x)$; equilibrium where $dU/dx = 0$
$U_s = \frac{1}{2}k(\Delta x)^2$; $U_g = -Gm_1 m_2 / r$; $\Delta U_g = mg\Delta y$	spring energy; gravity with zero at infinity (orbits); gravity near the surface
$P_{\text{avg}} = W/\Delta t = \Delta E/\Delta t$; $P_{\text{inst}} = dW/dt$	power ($P = F \cdot v$ is not printed)
$p = mv$; $F_{\text{net}} = dp/dt$; $J = \int F_{\text{net}} dt = \Delta p$; $v_{\text{cm}} = \Sigma p_i / \Sigma m_i$	momentum; variable mass or force; impulse (area under $F-t$); the centre of mass moves at constant velocity if no external force
Rotational (twin of)	Use when
$\omega = d\theta/dt$; $\alpha = d\omega/dt$; $\omega = \omega_0 + \alpha t$; $\theta = \theta_0 + \omega_0 t + \frac{1}{2}\alpha t^2$; $\omega^2 = \omega_0^2 + 2\alpha(\theta - \theta_0)$	angular kinematics (twins of v , a and the three equations)
$v = r\omega$; $a_T = r\alpha$; $\Delta x_{\text{cm}} = r\Delta\theta$	rim speed and tangential acceleration; rolling without slipping
$\tau = r \times F$; $\alpha_{\text{sys}} = \Sigma \tau / I_{\text{sys}} = \tau_{\text{net}} / I_{\text{sys}}$	torque ($rF \sin \theta$); Newton's second law for rotation (twin of $a = F/m$)
$I_{\text{tot}} = \Sigma I_i = \Sigma m_i r_i^2$; $I = \int r^2 dm$; $I' = I_{\text{cm}} + Md^2$	adding inertias; deriving I for a rod or disc; the parallel-axis theorem (rod about its end: $1/12 + 1/4 = 1/3 ML^2$)
$K_{\text{rot}} = \frac{1}{2}I\omega^2$; $W = \int \tau \cdot d\theta$	rotational energy (rolling: $\frac{1}{2}mv^2 + \frac{1}{2}I\omega^2$); rotational work
$L = r \times p = I\omega$; $\Delta L = \int \tau dt$	angular momentum (conserved when net torque is zero: skaters, bullet into a door); angular impulse
$T = 2\pi/\omega = 1/f$; $T_s = 2\pi\sqrt{(m/k)}$; $T_p = 2\pi\sqrt{(\ell/g)}$; $T_{\text{phys}} = 2\pi\sqrt{(I/(mgd))}$	periods: any oscillation; mass on a spring; simple pendulum (small angles); physical pendulum, d = pivot to centre of mass
$x = x_{\text{max}} \cos(\omega t + \phi)$	SHM position; differentiate for v and a (not printed): $v = -\omega x_{\text{max}} \sin(\omega t + \phi)$, $a = -\omega^2 x$

From

sciencewithshobha.com/blog/ap-physics-c-equation-sheet-explained-every-equation-on-the-official-mechanics-sheet-annotated-what-each-symbol-means-when-to-use-it-and-what-is-not-on-the-sheet/ · 1-on-1 online science and AP tutoring, KS2 to A level, Grades 5 to 12, free 30-minute first class.

The tables, what is NOT on the sheet, and a practice set

Table	Contents
Geometry and trig	$A = bh$, $\frac{1}{2}bh$, πr^2 ; $C = 2\pi r$; $s = r\theta$; $V = lwh$, $\pi r^2 l$ ($S = 2\pi r l + 2\pi r^2$), $\frac{4}{3}\pi r^3$ ($S = 4\pi r^2$); $a^2 + b^2 = c^2$, $\sin$, $\cos$, $\tan$
Vectors	$A \cdot B = AB \cos \theta$; $ A \times B = AB \sin \theta$; components $A_i + B_j + C_k$; $C = A + B$ by components
Calculus	chain rule; d/dx of x^n , $e^{\lambda(ax)}$, $\ln(ax)$, $\sin(ax)$, $\cos(ax)$; integrals of x^n ($n \neq -1$), $e^{\lambda(ax)}$, $1/(x+a)$, $\cos(ax)$, $\sin(ax)$
Identities	$\log(a \cdot b^x) = \log a + x \log b$; $\sin^2 \theta + \cos^2 \theta = 1$; $\sin 2\theta = 2 \sin \theta \cos \theta$; $\tan \theta = \sin \theta / \cos \theta$

Not on the sheet: know these

Missing	Know
Newton's laws in words	inertia; $F = ma$ as the sum of EXTERNAL forces; action-reaction pairs act on different bodies
Standard rotational inertias	rod about centre $\frac{1}{12} ML^2$, about end $\frac{1}{3} ML^2$; solid disc or cylinder $\frac{1}{2} MR^2$; hoop MR^2 ; solid sphere $\frac{2}{5} MR^2$; hollow sphere $\frac{2}{3} MR^2$
Rolling without slipping	$v_{cm} = r\omega$, $a_{cm} = r\alpha$; $K = \frac{1}{2}mv^2 + \frac{1}{2}I\omega^2$; friction does no work; $a = g \sin \theta / (1 + I/mr^2)$ down a ramp
Orbits	circular speed $\sqrt{GM/r}$; $K = \frac{1}{2}GMm/r$; total energy $-GMm/2r$; escape speed $\sqrt{2GM/r}$; Kepler's third law $T^2 = 4\pi^2 r^3 / GM$
Drag and terminal velocity	$F = -bv$ or $-cv^2$; terminal when drag = weight (mg/b); solve $m dv/dt = mg - bv$
Static equilibrium	$\Sigma F = 0$ and $\Sigma \tau = 0$ about any point
Collisions	momentum always conserved; K only in elastic; 1-D elastic: relative speed reverses
SHM derivatives; projectiles	$v = -\omega x_{max} \sin(\omega t + \phi)$, $a = -\omega^2 x$, $v_{max} = \omega x_{max}$; range $R = v^2 \sin 2\theta / g$
The derivations	where every sheet equation comes from: 'derive an expression' is the commonest free-response verb

Exam (2025-26): Section I 42 multiple choice, 85 min, 50%; Section II 4 free response, 95 min, 50%; hybrid digital (MCQ in Bluebook, FRQ handwritten); calculator allowed throughout; the sheet in both sections. The College Board has announced changes to the MCQ count and timings from May 2027: check the course page. Use the sheet to CHECK a sign or exponent, not to FIND an approach.

Practice: which sheet line, and what is missing?

- A rod of length L and mass M pivots about one end. Which sheet lines give its rotational inertia, and what must you supply?
- A satellite orbits at radius r . Which sheet line gives the force, and which missing fact gives the speed?
- A force $F(x) = 3x^2$ acts from $x = 0$ to 2 m. Which line gives the work, and what is it?
- A falling object has drag $F = -bv$. Which line sets up the equation of motion, and what is terminal velocity?
- A skater pulls her arms in. Which line explains why she spins faster?
- $U(x) = 4x^2 - 2x$. Which line gives the force, and where is equilibrium?

Answers

- $I = \int r^2 dm$ with $\lambda = M/L$ gives $\frac{1}{3} ML^2$; or $I = I_{cm} + Md^2$ with the missing $I_{cm} = \frac{1}{12} ML^2$ and $d = L/2$. 2. $|F_g| = Gm_1m_2/r^2$ with $a_c = v^2/r$ gives $v = \sqrt{GM/r}$, which is not printed. 3. $W = \int F \cdot dr = \int 3x^2 dx$ from 0 to $2 = x^3$ from 0 to $2 = 8$ J. 4. $F_{net} = dp/dt$ ($m dv/dt = mg - bv$); terminal when $mg = bv$, $v_t = mg/b$. 5. $L = I\omega$ with $\Delta L = \int \tau dt = 0$: I falls, so ω rises. 6. $F_x = -dU/dx = -(8x - 2)$; equilibrium where $8x - 2 = 0$, $x = 0.25$ m (stable: U is a minimum).